

***Additional Resources:***

1. *Relieving Pain in America: A Blueprint for Transforming Prevention, Care, Education, and Research*. National Academies Press; 2011. Accessed September 20, 2023. <https://nap.nationalacademies.org/catalog/13172/relieving-pain-in-america-a-blueprint-for-transforming-prevention-care>
2. *State of MSK Report 2021 Changing Musculoskeletal Trends Are Redefining MSK Care State of MSK Report 2021: Changing Musculoskeletal Trends Are Redefining MSK Care 2*. Accessed September 20, 2023. <https://healthactioncouncil.org/getmedia/a738c3c5-7c23-4739-bb8d-069dd5f7406b/Hinge-Health-State-of-MSK-Report-2021.pdf>
3. Jain A, Jain P, Aggarwal S. SARS-CoV-2 Impact on Elective Orthopaedic Surgery: Implications for Post-Pandemic Recovery. *J Bone Joint Surg Am*. 2020;102(13):e68. doi:10.2106/JBJS.20.00602
4. Demou E, Smith S, Bhaskar A, Mackay DF, Brown J, Hunt K, Vargas-Prada S, Macdonald EB. Evaluating sickness absence duration by musculoskeletal and mental health issues: a retrospective cohort study of Scottish healthcare workers. *BMJ Open*. 2018 Jan 26;8(1):e018085. doi: 10.1136/bmjopen-2017-018085. PMID: 29374662; PMCID: PMC5829784.
5. Liu X, Hanney WJ, Masaracchio M, et al. Immediate physical therapy initiation in patients with acute low back pain is associated with a reduction in downstream health care utilization and costs. *Phys Ther*. 2018;98(5):336-347.
6. Matifat E, Méquignon M, Cunningham C, Blake C, Fennelly O, Desmeules F. Benefits of musculoskeletal physical therapy in emergency departments: A systematic review. *Phys Ther*. 2019;99(9):1150-1166.
7. *J ortho Sports Phys Ther*, Vol 35, No 2, Feb 2005
8. Gatchel RJ, Polatin PB, Noe C, Gardea M, Pulliam C, Thompson J. Treatment- and cost-effectiveness of early intervention for acute low-back pain patients: a one-year prospective study. *J Occup Rehabil*. 2003;13(1):1-9.
9. Hahne AJ, Ford JJ, Surkitt LD, et al. Individualized physical therapy Is cost-effective compared with guideline-based advice for people with low back disorders. *Spine*. 2017;42(3):E169-E176.
10. Jensen BI, et al. A 3-year follow-up of a multidisciplinary rehabilitation programme for back and neck pain. *Pain*. 2005;115(3):273-283.

11. Awang H, et al. Factors related to successful return to work following multidisciplinary rehabilitation. *J Rehabil Med.* 2017;49(6):520-520.
12. Ojha HA, Wyrsta NJ, Davenport TE, Egan WE, Gellhorn AC. Timing of Physical Therapy Initiation for Nonsurgical Management of Musculoskeletal Disorders and Effects on Patient Outcomes: A Systematic Review. *J Orthop Sports Phys Ther.* 2016;46(2):56-70.
13. Liu X, Hanney WJ, Masaracchio M, et al. Immediate physical therapy initiation in patients with acute low back pain is associated with a reduction in downstream health care utilization and costs. *Phys Ther.* 2018;98(5):336-347.
14. Bürge E, et al. Cost-effectiveness of physical therapy only and of usual care for various health conditions: Systematic review. *Phys Ther.* 2016.
15. Childs JD, Fritz JM, Wu SS, et al. Implications of early and guideline adherent physical therapy for low back pain on utilization and costs. *BMC Health Serv Res.* 2015;15(1).
16. Fritz JM, Childs JD, Wainner RS, Flynn TW. Primary care referral of patients with low back pain to physical therapy. *Spine.* 2012;37(25):2114-2121.
17. Fritz JM, Brennan GP, Hunter SJ. Physical therapy or advanced imaging as first management strategy following a new consultation for low back pain in primary care: Associations with future health care utilization and charges. *Health Serv Res.* 2015;50(6):1927-1940.
18. Fritz JM, Kim M, Magel JS, Asche CV. Cost-effectiveness of primary care management with or without early physical therapy for acute low back pain: Economic evaluation of a randomized clinical trial. *Spine (Phila Pa 1976).* 2017;42(5):285-290.
19. Frogner BK, Harwood K, Andrilla CHA, Schwartz M, Pines JM. Physical therapy as the first point of care to treat low back pain: An instrumental variables approach to estimate impact on opioid prescription, health care utilization, and costs. *Health Serv Res.* 2018;53(6):4629-4646.
20. Miyamoto GC, Lin CC, Cabral CMN, van Dongen JM, van Tulder MW. Cost-effectiveness of exercise therapy in the treatment of non-specific neck pain and low back pain: a systematic review with meta-analysis. *Br J Sports Med.* 2019;53(3):172-181.
21. Neilson AR, Jones GT, Macfarlane GJ, et al. Cost-utility of maintained physical activity and physiotherapy in the management of distal arm pain: an economic evaluation of data from a randomized controlled trial. *Fam Pract.* 2019;36(2):179-186.

22. Peterson LE, Goodman C, et al. Assessment of the quality of cost analysis literature in physical therapy. *Phys Ther.* 2009;89(8):733-755.
23. Ojha HA, Wyrsta NJ, Davenport TE, Egan WE, Gellhorn AC. Timing of Physical Therapy Initiation for Nonsurgical Management of Musculoskeletal Disorders and Effects on Patient Outcomes: A Systematic Review. *J Orthop Sports Phys Ther.* 2016;46(2):56-70.
24. Andersen LN, Juul-Kristensen B, Sørensen TL, Herborg LG, Roessler KK, Søgaard K. Longer term follow-up of the effects of tailored physical activity or chronic pain self-management programme on return-to-work: a randomized controlled trial. *J Rehabil Med.* 2016;48(10):887-892.
25. Brendbekken R et al. Predictors of return to work in patients on sick leave with chronic musculoskeletal pain. *J Rehabil Med.* 2018;50(2).
26. Rivano M, Persson E, Stålnacke B-M, Schult M-L, Monika L. Return to work: One and two years follow-up after interdisciplinary pain rehabilitation. A study based on the Swedish Quality Registry for Pain Rehabilitation. *J Rehabil Med.* 2019;51(4):281-289.
27. Choi G, Raiturker PP, Kim M-J, Chung DJ, Chae Y-S, Lee S-H. The Effect of Early Isolated Lumbar Extension Exercise Program for Patients with Herniated Disc Undergoing Lumbar Discectomy. *Neurosurgery.* 2005;57(4):764-772.
28. Ozkara GO, Ozgen M, Ozkara E, Armagan O, Arslantas A, Atasoy MA. Effectiveness of physical therapy and rehabilitation programs starting immediately after lumbar disc surgery. *Turk Neurosurg.* 2015;25(3):372-379.
29. Fernandez-de-Las-Penas C, Cleland J, et al. The effectiveness of manual therapy versus surgery on self-reported function, cervical range of motion, and pinch grip force in carpal tunnel syndrome: A randomized clinical trial. *J Orthop Sports Phys Ther.* 2017;47(3):151-161.
30. Beissner KL, Saunders RL, McManis BG. Factors related to successful work hardening outcomes. *Phys Ther.* 1996;76(11):1188-1201.
31. Marin TJ, Van Eerd D, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain. *Cochrane Database Syst Rev.* 2017;6:CD002193.
32. Nieuwenhuijsen K, Faber B, Verbeek JH, et al. Interventions to improve return to work in depressed people. *Cochrane Database Syst Rev.* 2014.

33. Thijs K, Boer A, Vreugdenhil G, Wouw A, Houterman S, Schep G. Rehabilitation Using High-Intensity Physical Training and Long-Term Return-to-Work in Cancer Survivors. *J Occup Rehabil.* 2012;22(2):220-229.
34. Lechner D. Work hardening and work conditioning interventions: Do they affect disability? *Phys Ther.* 1994;74(5):471-493.
35. Niemeyer LO, Jacobs K, Reynolds-Lynch K, Bettencourt C, Lang S. Work Hardening: Past, present, and future -- The Work Programs Special Interest Section National Work-Hardening Outcome Study. *Am J Occup Ther.* 1994;48(4):327-339. 35.
36. Sang LS, Ying Eria LP. Outcome evaluation of work hardening program for manual workers with work-related back injury. *Work.* 2005;25(4):297-305.
37. Shin D. Characteristics of musculoskeletal disorders and satisfaction with in-house physical therapy clinics in office workers. *Work.* 2019;63(3):369-374.
38. Westgaard RH, Winkel J. Ergonomic intervention research for improved musculoskeletal health: A critical review. *Int J Ind Ergon.* 1997;20(6):463-500.
39. Andronis L, Kinghorn P, Qiao S, Whitehurst DGT, Durrell S, McLeod H. Cost-effectiveness of non-invasive and non-pharmacological interventions for low back pain: A systematic literature review. *Appl Health Econ Health Policy.* 2017;15(2):173-201.